





Nausea Prevention and Treatment

High: Aprepitant (IV)-Palonosetron-Dexamethasone-Lorazepam

- The following is a calendar to instruct you when to take your medicine.
- These medicines were prescribed to both prevent you from feeling nauseous ("sick to your stomach") and treat nausea if it occurs after your chemotherapy.
- The days and dates are listed down the left column. The <u>approximate</u> time of day that you should take the medicine is listed across the top.
- It is important to take your medicine as instructed <u>even if you are feeling well.</u> It is easier to prevent nausea and vomiting than to treat it once it occurs.
- If these medicines are not helping you or you think they are causing a side effect(s) please call 732-235-2465 and ask for the Nurse Help Line. Ask for your doctor if after 5 p.m. or on a weekend.

Aprepitant (IV)= Cinvanti[®], Palonosetron = Aloxi[®], Dexamethasone = Decadron[®], Lorazepam = Ativan[®]

Date	Medication	8 A.M.	8 P.M.
Day 1	Aprepitant (IV) 130 mg IV injection	Given by a nurse before chemotherapy	
Date:// Please Circle	Palonosetron 0.25 mg IV injection	Given by a nurse before chemotherapy	
Sn, M, T, W, Th, F, S	Dexamethasone 4 mg tablet	Given by a nurse before chemotherapy	
	Lorazepam 0.5 mg tablet		One (1) Tablet Only If Needed
Day 2	Dexamethasone	Two (2) Tablets	
Date://	4 mg tablet	with food or milk	
Please Circle Sn, M, T, W, Th, F, S	Lorazepam 0.5 mg tablet		One (1) Tablet Only If Needed
Day 3 Date://	Dexamethasone 4 mg tablet	Two (2) Tablets with food or milk	
Please Circle Sn, M, T, W, Th, F, S	Lorazepam 0.5 mg tablet		One (1) Tablet Only If Needed
Day 4	Dexamethasone	Two (2) Tablets	
Date://	4 mg tablet	with food or milk	
Please Circle Sn, M, T, W, Th, F, S	Lorazepam 0.5 mg tablet		One (1) Tablet Only If Needed

If you are in need of immediate assistance, please call 732-235-2465 and select the option that best meets your needs.